

Σύντομο Βιογραφικό Σημείωμα

Έμμου Ρογκάκου, Δρ.
Κλινική Επιγενετική και Μοριακή Ιατρική
Ερευνήτρια Β' βαθμίδας,

Ερευνητικό Πανεπιστημιακό Ινστιτούτο Υγείας Μητέρας Παιδιού και Ιατρικής Ακριβείας
Εθνικό και Καποδιστριακό Πανεπιστήμιο Αθηνών, Ιατρική Σχολή,
Α' Τμήμα Παιδιατρικής, Πανεπιστημιακό Νοσοκομείο Παιδών «Αγία Σοφία»,

Οδός: Θηβών και Παπαδιαμαντοπούλου,
11527, Αθήνα,
Ελλάδα.

e-mail: emmy.rogakou@med.uoa.gr

Tel: +30-2132013244
+30-6948207533

Ευρεσιτεχνίες

1. WO Patent App. PCT/US2000/017,288 - Method for determining DNA double-stranded breaks.
2. US Patent 6,884,873 - Kit for determining DNA double-stranded breaks with anti-γ-H2A antibodies.
3. US Patent 6,362,317 - Anti-γ-H2A antibody, fusion proteins thereof and method and kit for determining DNA double-stranded breaks.
4. National Patent (Greece), OBI (Hellenic Industrial Property Organisation)
5. National Patent (Greece), OBI (Hellenic Industrial Property Organisation)

Βιβλιογραφία

1. E.P.Rogakou, and K.E.Sekeri-Pataryas. (1993). Biochemical marker for differentiation is present in an *in vitro* aging cell system. **Biochem. Biophys. Res. Commun.** 196, 1274-1279.
2. E.P.Rogakou, D.R.Pilch, A.H.Orr, V.S.Ivanova, and W.M.Bonner. (1998). DNA double-strand breaks induce histone H2AX phosphorylation at serine-139. **J. Biol. Chem.** 273, 5858-5868.
3. E.P.Rogakou, C.Boon, C.Redon, W.M.Bonner. (1999) Megabase Chromatin Domains involved in DNA double-strand breaks *in vivo*. **J. Cell Biol.** 146, 905-916.
4. E.P.Rogakou, and K.E.Sekeri-Pataryas. (1999). Histone variants of H2A and H3 families are regulated during *in vitro* aging in the same manner as during differentiation. **Exp. Gerontology** 34, 741-754.
5. Futagi, Y., Suzuki, Y., Goto, M., Kato, T., Rogakou, E. P., & Sekeri-Pataryas, K. E. (1999). Histone variants of H2A and H3 families are regulated during *in vitro* aging in the same manner as during differentiation- Effects of aging and hydrocortisone. **Experimental Gerontology**, 34(6), 741-754.

6. E.P.Rogakou*, W.M.Bonner. (2000) Rapid histone extraction and electrophoretic analysis. *Biotechniques* 28, 38-46.
7. E.P.Rogakou*, W.Nieves-Neira, C.Boon, Y.Pommier, W.M.Bonner. (2000). Initiation of DNA fragmentation during apoptosis induces phosphorylation of H2AX histone at serine 139. *J. Biol. Chem.* 275, 9390-9395.
8. T.T.Paull, E.P.Rogakou, V.Yamazaki, C.U.Kirchgessner, M.Gellert, and W.M.Bonner. (2000) A critical role of histone H2AX in sequential recruitment of repair factors to nuclear foci after DNA damage. *Curr. Biol.* 10, 886-895.
9. H.T. Chen, A. Bhandoola, M.J. Difilippantonio, J. Zhu, M.J. Brown, X. Tai, E.P. Rogakou, T.M. Brotz, W.M. Bonner, T. Ried, A. Nussenzweig. (2000). Response to RAG-mediated V(D)J cleavage by NBS1 and γ -H2AX. *Science* 290, 1962-1965.
10. K. Johnson, W.M. Bonner, E.P. Rogakou, C. Redon, C Boon. (2000). Rapid histone extraction for electrophoretic analysis. *Biotechniques* 28, 38-46.
11. S.K.Mahadevaian, J.M. Turner, F.Baudat, E.P.Rogakou, P. de Boer, J.Blanco-Rodríguez, M.Jasin, S. Keeney, W.M.Bonner, and P.S.Burgoyne. (2001) Recombinational DNA double-strand breaks in mice precede synapsis. *Nature Genet.* 27, 271-276.
12. C.Redon, D.Pilch, E.Rogakou, O.Sedelnikova, K.Newrock, and W.M.Bonner. (2002) Histone H2A Variants H2AX and H2AZ. *Curr. Opin. Gen. Dev.* 12, 162-169.
13. O.A.Sedelnikova, E.P.Rogakou, I.G.Panyutin, W.M.Bonner. (2002) Quantitative detection of ¹²⁵JdU-induced DNA double-strand breaks with γ -H2AX antibody. *Rad. Research* 158, 486-492.
14. T. Furuta, Takemura H, Liao ZY, Aune GJ, Redon C, Sedelnikova OA, Pilch DR, Rogakou EP, Celeste A, Chen HT, Nussenzweig A, Aladjem MI, Bonner WM, Pommier Y. (2003) Phosphorylation of histone H2AX and activation of Mre11, Rad50, and Nbs1 in response to replication-dependent DNA double-strand breaks induced by mammalian DNA topoisomerase I cleavage complexes. *J.Biol.Chem.* 278, 20303-12.
15. C. Redon, D.R. Pilch, E.P. Rogakou, A.H. Orr, N.F. Lowndes, W.M. Bonner. (2003) Yeast histone 2A serine 129 is essential for the efficient repair of checkpoint-blind DNA damage. *EMBO Rep.* 4, 678-684.
16. R. Daniel, J. Ramcharan, E.P. Rogakou, K.D. Taganov, J.G. Greger, W.M. Bonner, A. Nussenzweig, R.A. Katz, A.M. Skalka (2004). Histone H2AX is phosphorylated at sites of retroviral DNA integration but is dispensable for postintegration repair, *J.Biol.Chem.* 279, 45810-45814.
17. I.V. Mavragani, Z. Nikitaki, M.P. Souli, A. Aziz, S. Nowsheen, K. Aziz, E.P. Rogakou, A.G. Georgakilas. (2017) Complex DNA damage: A route to radiation-induced genomic instability and carcinogenesis. *Cancers (Basel).* , 9(7), 91.
18. A. Georgoulis, C.E. Vorgias, G.P. Chrousos, E.P. Rogakou* (2017) Genome Instability and γ H2AX. *Int. J. Mol. Sci.* 18 (9), 1979-89.

(*corresponding author)

Κεφάλαια σε επιστημονικά βιβλία

1. Balatsos NAA, Rogakou EP*. (2005) “Histone H2AX involvement in DNA double-strand break repair pathways”, In “DNA Damage Recognition”. Siede W, New York, **Marcel Dekker Inc.** (*corresponding author).
2. Fatouros C., Rogakou EP*. (2007) “Interplay between gamma-H2AX and 53BP1 in DNA double-strand break repair pathways and apoptosis”, In “Apoptosis and Senescence in Cancer Chemotherapy and Radiotherapy”. Gewirtz D. A., New Jersey, **Humana Press**, (*corresponding author).
3. Rogakou EP*, Papadakis V., Chrousos GP. (2016) “The Epigenetic Biomarker γ H2AX: From Bench to Clinical Trials”. In “Epigenetics, the Environment, and Children’s Health Across Lifespans”. David Hollar, Switzerland, **Springer International Publishing** (*corresponding author).